

Some sufficient conditions on Hamiltonian digraph

Guohun Zhu

*Guilin University of Electronic Technology
No.1 Jinji Road, Guilin, China, 541004*

Abstract

Z-mapping graph is a balanced bipartite graph G of a digraph D by split each vertex of D into a pair of vertices of G . Based on the property of the G , it is proved that if D is strong connected and G is Hamiltonian, then D is Hamiltonian. It is also proved if D is Hamiltonian, then G contains at least a perfect matching. Thus some existence sufficient conditions for Hamiltonian digraph and Hamiltonian graph are proved to be equivalent, and two sufficient conditions of disjoint Hamiltonian digraph are given in this paper.

Key words: Z-mapping, Hamiltonian cycle, balanced bipartite graph, digraph

1 Introduction

It is well know that sufficient conditions for Hamiltonian cycle is study in graph always precede the digraph [Jorgen]. For instance the famous Dirac condition is presented in 1952.

Theorem 1 [Dirac] *Let G be a graph with $n > 2$ vertices. If $d(u) \geq n/2$ for every vertices, then G is Hamiltonian.*

The similar sufficient condition for digraph is presented in 1960.

Theorem 2 [Ghouila] *Let D be a strong connected digraph with $n > 2$ vertices. If $d(u) \geq n$ for every vertices, then D is Hamiltonian.*

It is not difficult to find that the degree in digraph is large twice than in undirected graph. Thus, we can think that the sufficient conditions of Hamiltonian digraph and the Hamiltonian graph have some relationship.

In this paper, A relationship is discover by a mapping between digraph and graph, which is named as Z-mapping graph [Zhu1] . The first this term is called "projector graph" when restrict the digraph with degree bound two [Zhu]. Based on this bijection, it is easy that some pervious sufficient conditions for Hamiltonian digraph and Hamiltonian graph are equivalent in the degree. And two disjoint Hamiltonian digraph is easy determined from the disjoint perfect matching bipartite graph.

Throughout the paper, we stick to the graph terminology and notation as given in [Harary] and [Zhu1] and focus on only finite simple (un)directed graph, i.e. the graph has no multi-arcs and no self loops. However, it need to recapitulate the cycle definition in graph and digraph with matrix forms respectively.

Definition 3 *A simple cycle L in digraph is a set of arcs $(a_1, a_2, \dots, a_l)$, the incidence matrix C could be permutated by row or column exchange to following forms*

$$C = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & -1 \\ -1 & 1 & 0 & \dots & 0 & 0 \\ 0 & -1 & 1 & \dots & 0 & 0 \\ 0 & 0 & -1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & -1 & 1 \end{pmatrix}. \quad (1)$$

Definition 4 *A simple cycle L in graph is a set of edges $(e_1, e_2, \dots, e_l)$, the incidence matrix C could be permutate by row col exchange to following forms*

$$C = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 \\ 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 1 & 1 \end{pmatrix}. \quad (2)$$

The cycle definition responses to an edge set of G , which is easy to build a mapping from matching which is another definition on an edge set of graph G .

2 Z-mapping

Firstly, let us recapitulate the incidence matrix divided method in [Zhu1], which divided the incidence matrix C of a digraph into two groups.

$$C^+ = \{c_{ij} | c_{ij} \geq 0 \text{ otherwise } 0\} \quad (3)$$

$$C^- = \{c_{ij} | c_{ij} \leq 0 \text{ otherwise } 0\} \quad (4)$$

Secondly, let us combine these two matrices into a matrix.

Definition 5 Let C be a incidence matrix of digraph D , the Z -mapping graph of D , denoted as $Z(D)$, is defined as a balanced bipartite graph $G(X, Y : E)$

with a incidence matrix $F = \begin{pmatrix} C^+ \\ -C^- \end{pmatrix}$,

In fact, it is a isomorphism build from $D(V, A)$ to $G(X, Y; E)$. Let the $V^+ = \{v_i^+\}$ represents the vertices in C^+ and the $V^- = \{v_i^-\}$ represents the vertices in C^- . More precisely, the mapping $D(V, A) \rightarrow G(V^+, V^-; E)$ defines a map from arcs of D to edges of G :

$$E = \{(v_i^+, v_j^-) | \langle v_i, v_j \rangle \in A\}. \quad (5)$$

Based on the same principle, we also define a reverse mapping from a simple balance bipartite graph G to a simple digraph D .

$$f^{-1} : G(V^+, V^-; E) \rightarrow D(V; A) \quad (6)$$

this function is a reverse function of the equation 5:

$$A = \{\langle x, y \rangle | (x, y) \in E \wedge x \in V^+ \wedge y \in V^-\}. \quad (7)$$

Thus first main result is follows:

Theorem 6 Given a Z -mapping graph G of a strong connected digraph D , if G is Hamiltonian, then D is Hamiltonian.

PROOF. Suppose the G has a Hamiltonian cycle L and the response incidence matrix is a $2n \times 2n$ matrix C .

$$C = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 & 1 \\ 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 \\ 0 & 0 & 0 & \dots & 1 & 1 \end{pmatrix}. \quad (8)$$

And divide C to two groups: C_1 and C_2 , where

$$C_1 = \{c_i | i \text{ is odd}\}$$

$$C_2 = \{c_i | i \text{ is even}\}$$

Let $C' = C_1 - C_2$. Then obtain a matrix $n \times 2n$.

$$C' = \begin{pmatrix} 0 & -1 & 0 & \dots & 0 & 1 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & -1 \end{pmatrix}. \quad (9)$$

After the column from C' with all zero is removed. Then a new matrix $n \times n$ is obtained. This matrix satisfies the cycle equation 1.

The second main result is follows:

Theorem 7 *Given a Z-mapping graph G of a digraph D , if D contains a Hamiltonian cycle L , then G contains a perfect matching M , where $M = F(L)$.*

PROOF. Suppose that D has a Hamiltonian cycle L , then D has a sub incidence matrix as equation 1. According to the divided matrices approach, the response Z-mapping graph $G(X, Y; E)$ is a 1-regular bipartite graph, thus $\forall x \in X \ |N(x)| = |X|$, thus according to the Hall [Hall]rules, the graph z-mapping graph G is a perfect matching.

3 Sufficient conditions from Hamiltonian Graphs to Digraphs

According to the theorem 6, the theorem 2 can be directly generated from theorem 1. As the Dirac theorem has extended to the $d(u) < n/2$ case by the Faudree etc. as follows.

Theorem 8 [Faudree] *Let $G(V, E)$ be a graph with $n > 2$ vertices and a subset $S = \{v : d(v) < \frac{n}{2}\}$ of V , if k is the minimum degree of G , and $|S| \leq k - 1$ then G is Hamiltonian.*

Thus as the theorem 6, it can be derived to a new theorem.

Theorem 9 *Let D be a strong connected digraph with $n > 2$ vertices and $S = \{v : d(v) < n\}$. if k is the minimum degree of D , and $|S| \leq k - 1$, then D is Hamiltonian.*

In fact, Moon and Moser [Moon] have presents a theorem for similar sufficient condition on bipartite Hamiltonian graph.

Theorem 10 [Moon] *Let B be $2n$ balance bipartite graph, where $1 \leq k \leq n$, if the number of vertices $S = \{v : d(v) < k\}$ $k < n$ is less than n , then B is Hamiltonian.*

In [Moon], Moon and Moser also presented a theorem for balanced bipartite graph as follows.

Theorem 11 [Moon] *Given a balanced bipartite graph with $2n > 2$ vertices, for all vertices if $d(u) > \frac{n}{2}$, then G is Hamiltonian.*

It is easy to find that the theorem 11 could derive a sufficient condition $d^+(u) + d^-(u) = d(u) > n$ for Hamiltonian digraph, but this is not precede the theorem 2.

Since a Hamiltonian cycle consists of two independent perfect matching in balanced bipartite graph. Thus a sufficient condition for Hamiltonian digraph as follows.

COROLLARY 1 *Given a strong connected digraph with $n > 2$ vertices. for all vertices if $d^+(u) > \frac{n}{2}$ and $d^-(u) > \frac{n}{2}$, then D contains two disjoint Hamiltonian cycle.*

4 Sufficient conditions from Hamiltonian digraph to perfect bipartite graph

The relation perfect matching and cycle . For instance, in 1972, Las Verganas obtained the following results:

Theorem 12 [Verganas] *Let $G(X, Y; E)$ be a balanced bipartite graph with $2n > 2$ vertices. For all vertices $u \in X$, $v \in Y$, $(u, v) \notin E$, If $d(u) + d(v) \geq n + 2$, then every perfect matching in G is contained in a Hamiltonian cycle.*

According to theorem 6, it could derived a sufficient condition for Hamiltonian digraph from Ore conditions [Ore] such as $d(u) + d(v) \geq n$, but Woodall had obtained a more strict sufficient condition.

Theorem 13 [Woodall] *Let D be a strong connected digraph with $n > 2$ vertices. For all vertices u, v , $\langle u, v \rangle \notin A$, If $d^+(u) + d^-(v) \geq n$, then D is Hamiltonian.*

Combination these two theorems, we could get two interesting results.

COROLLARY 2 *Let $D(V, A)$ be a digraph with $n > 2$ vertices. For all vertices u, v , $\langle u, v \rangle \notin A$, If $d^+(u) + d^-(v) \geq n + 2$, then D contains at least two disjoint Hamiltonian cycles.*

COROLLARY 3 *Let $G(X, Y; E)$ be a balanced bipartite graph with $2n > 2$ vertices. For all vertices $u \in X$, $v \in Y$, $(u, v) \notin E$, If $d(u) + d(v) \geq n$, then G contains at least one perfect matching, if $d(u) + d(v) \geq n + 2$, then G contains at least two disjoint perfect matchings.*

References

- [Jorgen] J. Bang-Jensen. Yubao Guo, Anders Yeo. A new sufficient condition for a digraph to be Hamiltonian. Discrete Applied Mathematics. (1999) 95:61-72.
- [Dirac] G. A. Dirac, Some theorems on abstract graphs. In Proceedings of the London Mathematical Society, third series, Vol.2, March 1952, pp.69–81.
- [Ghouila] A. Ghouila-Houri, Une condition suffisante d'existence d'un circuit hamiltonien, CR Acad. Sci. Paris 25 (1960) 495–497.
- [Faudree] R. J. Faudree, Richard H. Schelp, Akira Saito, Ingo Schiermeyer. Degree conditions for Hamiltonicity: Counting the number of missing edges. Discrete Mathematics 307 (2007) 873–877.
- [Harary] F. Harary, Graph Theory. Addison-Wesley Series in Mathematics. (1969). 150–160.

- [Hall] P. HALL, On representatives of subsets, J. London Math. Soc. 10 (1936), 26–30.
- [Moon] J. Moon, L. Moser, On hamiltonian bipartite graphs, Israel J. Math. 1 (1963) 163–165.
- [Ore] O. Ore, Note on Hamiltonian circuits. Am Math Monthly 67 (1960), 55.
- [Verganas] M. L. Vergnas, Problèmes de couplages et problemes hamiltoniens en théorie des graphes, Ph.D. Thesis, Université Paris Xi, 1972.
- [Woodall] D.R. Woodall. Sufficient conditions for cycles in digraphs, Proc. London Math. Soc. 24(1972), 739–755.
- [Zhu] G. Zhu, The Complexity of Hamiltonian Cycle Problem in Digraphs with Degree Bound Two is Polynomial Time, arXiv:0704.0309v2, 2007.
- [Zhu1] G. Zhu, On the Complexity of Hamiltonian Cycle Problem in Digraphs with Degree Bound Two, Computational Intelligence and Industrial Applications: Proceedings of ISCIIA2008. (2008) 227-234.